

# Convex sets and convex functions

Mauro Passacantando

Department of Computer Science, University of Pisa  
Largo Pontecorvo 3  
mauro.passacantando@unipi.it

Optimization Methods  
Master of Science in Embedded Computing Systems – University of Pisa  
<http://pages.di.unipi.it/passacantando/om/OM.html>

## Subspaces

Given  $x, y \in \mathbb{R}^n$ .

A **linear combination** of  $x$  and  $y$  is a point  $\alpha x + \beta y$ , where  $\alpha, \beta \in \mathbb{R}$ .

A set  $C \subseteq \mathbb{R}^n$  is a **subspace** if it contains all the linear combinations of any two points in  $C$ .

### Examples:

- ▶  $\{0\}$
- ▶ any line which passes through zero
- ▶ the solution set of a homogeneous system of linear equations

$$C = \{x \in \mathbb{R}^n : Ax = 0\},$$

where  $A$  is a  $m \times n$  matrix.

## Affine sets

An **affine combination** of  $x$  and  $y$  is a point  $\alpha x + \beta y$ , where  $\alpha + \beta = 1$ .

A set  $C \subseteq \mathbb{R}^n$  is an **affine** set if it contains all the affine combinations of any two points in  $C$ .

### Examples:

- ▶ any single point  $\{x\}$
- ▶ any line
- ▶ the solution set of a system of linear equations

$$C = \{x \in \mathbb{R}^n : Ax = b\},$$


where  $A$  is a  $m \times n$  matrix and  $b \in \mathbb{R}^m$

- ▶ any subspace


## Convex sets

A **convex combination** of two given points  $x$  and  $y$  is a point  $\alpha x + \beta y$ , where  $\alpha + \beta = 1$ ,  $\alpha \geq 0$ ,  $\beta \geq 0$ .

A set  $C \subseteq \mathbb{R}^n$  is **convex** if it contains all the convex combinations of any two points in  $C$ .



convex set




non-convex set

**Exercise.** Prove that if  $C$  is convex, then for any  $x^1, \dots, x^k \in C$  and  $\alpha_1, \dots, \alpha_k \in (0, 1)$  s.t.  $\sum_{i=1}^k \alpha_i = 1$ , one has  $\sum_{i=1}^k \alpha_i x^i \in C$ .

## Convex hull

The **convex hull**  $\text{conv}(C)$  of a set  $C$  is the smallest convex set containing  $C$ .



**Exercise.** Prove that  $\text{conv}(C) = \{\text{all convex combinations of points in } C\}$ .

**Exercise.** Prove that  $C$  is convex if and only if  $C = \text{conv}(C)$ .

## Convex sets - Examples

### Examples:

- ▶ subspace
- ▶ affine set
- ▶ line segment
- ▶ halfspace  $\{x \in \mathbb{R}^n : a^T x \leq b\}$
- ▶ polyhedron  $P = \{x \in \mathbb{R}^n : Ax \leq b\}$  solution set of a system of linear inequalities

## Convex sets - Examples

- ▶ ball  $B(x, r) = \{y \in \mathbb{R}^n : \|y - x\| \leq r\}$ , where  $\|\cdot\|$  is any norm, e.g.

$$\|x\|_2 = \sqrt{\sum_{i=1}^n x_i^2} \text{ (Euclidean norm)}$$

$$\|x\|_1 = \sum_{i=1}^n |x_i| \text{ (Manhattan distance)}$$

$$\|x\|_\infty = \max_{i=1, \dots, n} |x_i| \text{ (Chebyshev norm)}$$

$$\|x\|_p = \sqrt[p]{\sum_{i=1}^n |x_i|^p}, \text{ with } 1 \leq p \leq \infty$$

$$\|x\|_A = \sqrt{x^T A x}, \text{ where } A \text{ is a symmetric and positive definite matrix, i.e.,}$$

$$x^T A x > 0 \quad \forall x \neq 0.$$

**Exercise.** Find  $B(0, 1)$  w.r.t.  $\|\cdot\|_1$ ,  $\|\cdot\|_\infty$  and  $\|\cdot\|_A$  where  $A = \begin{pmatrix} 2 & 0 \\ 0 & 1 \end{pmatrix}$ .

## Operations that preserve convexity

### Sum and difference

If  $C_1$  and  $C_2$  are convex, then  $C_1 + C_2 := \{x + y : x \in C_1, y \in C_2\}$  is convex.

If  $C_1$  and  $C_2$  are convex, then  $C_1 - C_2 := \{x - y : x \in C_1, y \in C_2\}$  is convex.

### Intersection

If  $C_1$  and  $C_2$  are convex, then  $C_1 \cap C_2$  is convex.

**Exercise.** If  $\{C_i\}_{i \in I}$  is a family of convex sets, then  $\bigcap_{i \in I} C_i$  is convex.

### Union

If  $C_1$  and  $C_2$  are convex, then  $C_1 \cup C_2$  is convex?

### Closure and interior

If  $C$  is convex, then  $\text{cl}(C)$  is convex.

If  $C$  is convex, then  $\text{int}(C)$  is convex.



## Operations that preserve convexity

### Affine functions

Let  $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$  be affine, i.e.  $f(x) = Ax + b$ , with  $A \in \mathbb{R}^{m \times n}$ ,  $b \in \mathbb{R}^m$ .

- ▶ If  $C \subseteq \mathbb{R}^n$  is convex, then  $f(C) = \{f(x) : x \in C\}$  is convex
- ▶ If  $C \subseteq \mathbb{R}^m$  is convex, then  $f^{-1}(C) = \{x \in \mathbb{R}^n : f(x) \in C\}$  is convex

### Examples:

- ▶ scaling, e.g.  $f(x) = \alpha x$ , with  $\alpha > 0$
- ▶ translation, e.g.  $f(x) = x + b$ , with  $b \in \mathbb{R}^n$
- ▶ rotation, e.g.  $f(x) = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} x$ , with  $\theta \in (0, 2\pi)$

## Cones

A set  $C \subseteq \mathbb{R}^n$  is a **cone** if  $\alpha x \in C$  for any  $x \in C$  and  $\alpha \geq 0$ .

### Examples:

- ▶  $\mathbb{R}_+^n$  is a convex cone
- ▶  $\{x \in \mathbb{R}^2 : x_1 x_2 = 0\}$  is a nonconvex cone
- ▶ Given a polyhedron  $P = \{x : Ax \leq b\}$ , the recession cone of  $P$  is defined as

$$\text{rec}(P) := \{d : x + \alpha d \in P \text{ for any } x \in P, \alpha \geq 0\}.$$

It is easy to prove  $\text{rec}(P) = \{x : Ax \leq 0\}$ , thus it is a polyhedral cone.

- ▶  $\{x \in \mathbb{R}^3 : x_3 \geq \sqrt{x_1^2 + x_2^2}\}$  is a non-polyhedral cone.

## Exercises

- Write the vector  $(1, 1)$  as the convex combination of the vectors  $(0, 0)$ ,  $(3, 0)$ ,  $(0, 2)$ ,  $(3, 2)$ .
- When does one halfspace contain another? Give conditions under which

$$\{x \in \mathbb{R}^n : a_1^T x \leq b_1\} \subseteq \{x \in \mathbb{R}^n : a_2^T x \leq b_2\},$$


where  $\|a_1\|_2 = \|a_2\|_2 = 1$ . Also find the conditions under which the two halfspaces are equal.

- Which of the following sets are polyhedra?
  - $\{y_1 a_1 + y_2 a_2 : -1 \leq y_1 \leq 1, -1 \leq y_2 \leq 1\}$ , where  $a_1, a_2 \in \mathbb{R}^n$ .
  - $\left\{x \in \mathbb{R}^n : x \geq 0, \sum_{i=1}^n x_i = 1, \sum_{i=1}^n a_i x_i = b_1, \sum_{i=1}^n a_i^2 x_i = b_2\right\}$ , where  $b_1, b_2, a_1, \dots, a_n \in \mathbb{R}$ .
  - $\{x \in \mathbb{R}^n : x \geq 0, a^T x \leq 1 \text{ for all } a \text{ with } \|a\|_2 = 1\}$ .
  - $\{x \in \mathbb{R}^n : x \geq 0, a^T x \leq 1 \text{ for all } a \text{ with } \|a\|_1 = 1\}$ .

## Convex functions

Given a convex set  $C \subseteq \mathbb{R}^n$ , a function  $f : C \rightarrow \mathbb{R}$  is **convex** if

$$f(\alpha y + (1 - \alpha)x) \leq \alpha f(y) + (1 - \alpha)f(x) \quad \forall x, y \in C, \forall \alpha \in (0, 1)$$



$f$  is said **concave** if  $-f$  is convex.

**Exercise.** Prove that if  $f$  is convex, then for any  $x^1, \dots, x^k \in C$  and  $\alpha_1, \dots, \alpha_k \in (0, 1)$  s.t.  $\sum_{i=1}^k \alpha_i = 1$ , one has  $f\left(\sum_{i=1}^k \alpha_i x^i\right) \leq \sum_{i=1}^k \alpha_i f(x^i)$ .

## Strictly convex and strongly convex functions

Given a convex set  $C \subseteq \mathbb{R}^n$ , a function  $f : C \rightarrow \mathbb{R}$  is **strictly convex** if

$$f(\alpha y + (1 - \alpha)x) < \alpha f(y) + (1 - \alpha)f(x) \quad \forall x, y \in C, \forall \alpha \in (0, 1)$$

Given a convex set  $C \subseteq \mathbb{R}^n$ , a function  $f : C \rightarrow \mathbb{R}$  is **strongly convex** if there exists  $\tau > 0$  s.t.

$$f(\alpha y + (1 - \alpha)x) \leq \alpha f(y) + (1 - \alpha)f(x) - \frac{\tau}{2}\alpha(1 - \alpha)\|y - x\|^2 \\ \forall x, y \in C, \forall \alpha \in (0, 1)$$

**Thm.**  $f$  is strongly convex if and only if  $\exists \tau > 0$  s.t.  $f(x) - \frac{\tau}{2}\|x\|^2$  is convex

### Exercise.

- ▶ Prove that: strongly convex  $\implies$  strictly convex  $\implies$  convex
- ▶ convex  $\implies$  strictly convex ?
- ▶ strictly convex  $\implies$  strongly convex ?


## First order conditions

Assume that  $C \subseteq \mathbb{R}^n$  is open convex and  $f : C \rightarrow \mathbb{R}$  is continuously differentiable.

### Theorem

$f$  is **convex** if and only if

$$f(y) \geq f(x) + (y - x)^T \nabla f(x) \quad \forall x, y \in C.$$



First-order approximation of  $f$  is a global **underestimator**

## First order conditions

### Theorem

- ▶  $f$  is **strictly convex** if and only if

$$f(y) > f(x) + (y - x)^T \nabla f(x) \quad \forall x, y \in C, \text{ with } x \neq y.$$

- ▶  $f$  is **strongly convex** if and only if there exists  $\tau > 0$  such that

$$f(y) \geq f(x) + (y - x)^T \nabla f(x) + \frac{\tau}{2} \|y - x\|_2^2 \quad \forall x, y \in C.$$

## Second order conditions

Assume that  $C \subseteq \mathbb{R}^n$  is open convex and  $f : C \rightarrow \mathbb{R}$  is twice continuously differentiable.

### Theorem

- ▶  $f$  is **convex** if and only if for all  $x \in C$  the Hessian matrix  $\nabla^2 f(x)$  is positive semidefinite, i.e.

$$v^T \nabla^2 f(x) v \geq 0 \quad \forall v \neq 0,$$

or, equivalently, the eigenvalues of  $\nabla^2 f(x)$  are  $\geq 0$ .

- ▶ If  $\nabla^2 f(x)$  is positive definite for all  $x \in C$ , then  $f$  is **strictly convex**.
- ▶  $f$  is **strongly convex** if and only if there exists  $\tau > 0$  such that  $\nabla^2 f(x) - \tau I$  is positive semidefinite for all  $x \in C$ , i.e.

$$v^T \nabla^2 f(x) v \geq \tau \|v\|_2^2 \quad \forall v \neq 0,$$

or, equivalently, the eigenvalues of  $\nabla^2 f(x)$  are  $\geq \tau$ .



## Examples

$f(x) = c^T x$  is both convex and concave

$f(x) = \frac{1}{2}x^T Qx + c^T x$  is

- ▶ convex iff  $Q$  is positive semidefinite
- ▶ strongly convex iff  $Q$  is positive definite
- ▶ concave iff  $Q$  is negative semidefinite
- ▶ strongly concave iff  $Q$  is negative definite

$f(x) = e^{ax}$  for any  $a \in \mathbb{R}$  is strictly convex, but not strongly convex

$f(x) = \log(x)$  is strictly concave, but not strongly concave

$f(x) = x^a$  with  $x > 0$  is strictly convex if  $a > 1$  or  $a < 0$ . Is it strongly convex?

$f(x) = x^a$  with  $x > 0$  is strictly concave if  $0 < a < 1$

$f(x) = \|x\|$  is convex, but not strictly convex

$f(x) = \max\{x_1, \dots, x_n\}$  is convex, but not strictly convex

## Exercises

1. Prove that the function

$$f(x_1, x_2) = \frac{x_1 x_2}{x_1 - x_2}$$

is convex on the set  $\{x \in \mathbb{R}^2 : x_1 - x_2 > 0\}$ .

2. Prove that  $f(x_1, x_2) = \frac{1}{x_1 x_2}$  is convex on the set  $\{x \in \mathbb{R}^2 : x_1, x_2 > 0\}$ .
3. Given a convex set  $C \subseteq \mathbb{R}^n$ , the distance function is defined as follows:

$$d_C(x) = \inf_{y \in C} \|y - x\|.$$

Prove that  $d_C$  is a convex function.

4. Given  $C = \{x \in \mathbb{R}^2 : x_1^2 + x_2^2 \leq 1\}$ , write the distance function  $d_C$  explicitly.
5. Prove that the arithmetic mean of  $n$  positive numbers  $x_1, \dots, x_n$  is greater or equal to their geometric mean, i.e.,

$$\frac{x_1 + x_2 + \dots + x_n}{n} \geq \sqrt[n]{x_1 x_2 \dots x_n}.$$

(Hint: exploit the  $\log$  function.)

## Operations that preserve convexity

### Theorem

- ▶ If  $f$  is convex and  $\alpha > 0$ , then  $\alpha f$  is convex
- ▶ If  $f_1$  and  $f_2$  are convex, then  $f_1 + f_2$  are convex
- ▶ If  $f$  is convex, then  $f(Ax + b)$  is convex

### Examples

- ▶ Log barrier for linear inequalities:

$$f(x) = - \sum_{i=1}^m \log(b_i - a_i^T x) \quad C = \{x \in \mathbb{R}^n : b_i - a_i^T x > 0 \quad \forall i = 1, \dots, m\}$$

- ▶ Norm of affine function:  $f(x) = \|Ax + b\|$

**Exercise.** If  $f_1$  and  $f_2$  are convex, then is the product  $f_1 f_2$  convex?

## Pointwise maximum

### Theorem

- ▶ If  $f_1, \dots, f_m$  are convex, then  $f(x) = \max\{f_1(x), \dots, f_m(x)\}$  is convex.
- ▶ If  $\{f_i\}_{i \in I}$  is a family of convex functions, then  $f(x) = \sup_{i \in I} f_i(x)$  is convex.

**Example.** If  $L(x, \lambda) : \mathbb{R}^n \times \mathbb{R}^m \rightarrow \mathbb{R}$  is convex in  $x$  and concave in  $\lambda$ , then

$$\begin{aligned} p(x) &= \sup_{\lambda} L(x, \lambda) && \text{is convex} \\ d(\lambda) &= \inf_x L(x, \lambda) && \text{is concave} \end{aligned}$$

## Composition

$f : \mathbb{R}^n \rightarrow \mathbb{R}$  and  $g : \mathbb{R} \rightarrow \mathbb{R}$ .

### Theorem

- ▶ If  $f$  is convex and  $g$  is convex and nondecreasing, then  $g \circ f$  is convex.
- ▶ If  $f$  is concave and  $g$  is convex and nonincreasing, then  $g \circ f$  is convex.
  
- ▶ If  $f$  is concave and  $g$  is concave and nondecreasing, then  $g \circ f$  is concave.
- ▶ If  $f$  is convex and  $g$  is concave and nonincreasing, then  $g \circ f$  is concave.

### Examples

- ▶ If  $f$  is convex, then  $e^{f(x)}$  is convex
- ▶ If  $f$  is concave and positive, then  $\log f(x)$  is concave
- ▶ If  $f$  is convex, then  $-\log(-f(x))$  is convex on  $\{x : f(x) < 0\}$
- ▶ If  $f$  is concave and positive, then  $\frac{1}{f(x)}$  is convex
- ▶ If  $f$  is convex and nonnegative, then  $f(x)^p$  is convex for all  $p \geq 1$

## Sublevel sets

Given  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  and  $\alpha \in \mathbb{R}$ , the set

$$S_\alpha(f) = \{x \in \mathbb{R}^n : f(x) \leq \alpha\}$$

is said the  $\alpha$ -sublevel set of  $f$ .

**Exercise.** Prove that if  $f$  is convex, then  $S_\alpha(f)$  is a convex set for any  $\alpha \in \mathbb{R}$ .

Is the converse true?

## Quasiconvex functions

Given a convex set  $C \subseteq \mathbb{R}^n$ , a function  $f : C \rightarrow \mathbb{R}$  is **quasiconvex** if the  $\alpha$ -sublevel sets are convex for all  $\alpha \in \mathbb{R}$ .

$f$  is said quasiconcave if  $-f$  is quasiconvex.

### Examples

- ▶  $f(x) = \sqrt{|x|}$  is quasiconvex on  $\mathbb{R}$
- ▶  $f(x_1, x_2) = x_1 x_2$  is quasiconcave on  $\{x \in \mathbb{R}^2 : x_1 > 0, x_2 > 0\}$
- ▶  $f(x) = \log x$  is quasiconvex and quasiconcave
- ▶  $f(x) = \text{ceil}(x) = \inf\{z \in \mathbb{Z} : z \geq x\}$  is quasiconvex and quasiconcave

## Exercise

Express each convex set defined below in the form  $\bigcap_{i \in I} \{x : f_i(x) \leq 0\}$ , where  $f_i : \mathbb{R}^n \rightarrow \mathbb{R}$  are suitable convex functions:

**a)**  $\text{conv}\{(-1, -1), (1, 0), (0, 2)\}$

**b)**  $\text{conv}\{(0, 0), (1, 1)\}$

**c)**  $\text{conv}(\{x \in \mathbb{R}^2 : x_1^2 + (x_2 - 1)^2 = 1\} \cup \{x \in \mathbb{R}^2 : x_1^2 + (x_2 + 1)^2 = 1\})$

**d)**  $\text{conv}\{x \in \mathbb{R}^2 : x_1 x_2 = 1\}$