

# Solution methods for unconstrained optimization problems

Mauro Passacantando

Department of Computer Science, University of Pisa  
mauro.passacantando@unipi.it

Optimization Methods  
Master of Science in Embedded Computing Systems – University of Pisa  
<http://pages.di.unipi.it/passacantando/om/OM.html>

## Gradient method

Current point  $x^k$ , search direction  $d^k = -\nabla f(x^k)$  (steepest descent direction)

### Gradient method

Choose  $x^0 \in \mathbb{R}^n$ , set  $k = 0$

**while**  $\nabla f(x^k) \neq 0$  **do**

$$d^k = -\nabla f(x^k)$$

compute an optimal solution  $t_k$  of the problem:  $\min_{t>0} f(x^k + t d^k)$

$$x^{k+1} = x^k + t_k d^k, k = k + 1$$

**end**

**Example.**  $f(x) = x_1^2 + 2x_2^2 - 3x_1 - 2x_2$ , starting point  $x^0 = (2, 1)$ .

$$\nabla f(x^0) = (1, 2), d^0 = (-1, -2), f(x^0 + t d^0) = 9t^2 - 5t - 2, t_0 = 5/18,$$

$$x^1 = (2, 1) - \frac{5}{18}(1, 2) = \left(\frac{31}{18}, \frac{4}{9}\right).$$

## Gradient method - convergence

**Exercise 1.** Prove that  $(d^k)^T d^{k+1} = 0$  for any iteration  $k$ .

**Exercise 2.** Prove that if  $\{x^k\}$  converges to  $x^*$ , then  $\nabla f(x^*) = 0$ , i.e.  $x^*$  is a stationary point of  $f$ .

### Theorem

If  $f$  is **coercive**, then for any starting point  $x^0$  the generated sequence  $\{x^k\}$  is bounded and any of its cluster points is a **stationary point** of  $f$ .

### Corollary

If  $f$  is **coercive and convex**, then for any starting point  $x^0$  the generated sequence  $\{x^k\}$  is bounded and any of its cluster points is a **global minimum** of  $f$ .

### Corollary

If  $f$  is **strongly convex**, then for any starting point  $x^0$  the generated sequence  $\{x^k\}$  converges to the **global minimum** of  $f$ .

## Gradient method - step size

If  $f(x) = \frac{1}{2}x^T Qx + c^T x$ , with  $Q$  positive definite matrix, then

$$f(x^k + td^k) = f(x^k - tg^k) = \frac{1}{2} (g^k)^T Q g^k t^2 - (g^k)^T g^k t + f(x^k),$$

where  $g^k = \nabla f(x^k) = Qx^k + c$ . Thus the step size is

$$t_k = \frac{(g^k)^T g^k}{(g^k)^T Q g^k} = -\frac{(g^k)^T d^k}{(d^k)^T Q d^k}.$$

## Gradient method - exercises

**Exercise 3.** Implement in MATLAB the gradient method for solving the problem

$$\begin{cases} \min & \frac{1}{2}x^T Qx + c^T x \\ x \in & \mathbb{R}^n \end{cases}$$

where  $Q$  is a positive definite matrix.

**Exercise 4.** Run the gradient method for solving the problem

$$\begin{cases} \min & 3x_1^2 + 3x_2^2 + 3x_3^2 + 3x_4^2 - 4x_1x_3 - 4x_2x_4 + x_1 - x_2 + 2x_3 - 3x_4 \\ x \in & \mathbb{R}^4 \end{cases}$$

starting from the point  $(0, 0, 0, 0)$ . [Use  $\|\nabla f(x)\| < 10^{-6}$  as stopping criterion.]

**Exercise 5.** Consider the following data fitting problem: given the experimental data in fitting.txt, find the best approximating polynomial of degree 3 w.r.t. the Euclidean norm. Solve the problem by means of the gradient method starting from  $x^0 = 0$ . [Use  $\|\nabla f(x)\| < 10^{-3}$  as stopping criterion.]

## Gradient method - step size

If  $f$  is a general nonlinear function, how to find the step size  $t_k$ ?

Assume that the restriction  $\varphi(t) = f(x^k + td^k)$  is strongly convex, so we have to find  $t^*$  s.t.  $\varphi'(t^*) = 0$ .

### Bisection method

Find  $b > 0$  s.t.  $\varphi'(b) > 0$  (we know that  $\varphi'(0) < 0$ ).

$$a_0 := 0, b_0 := b, t_0 := \frac{a_0 + b_0}{2}, i := 0.$$

**while**  $\varphi'(t_i) \neq 0$  **do**

**if**  $\varphi'(t_i) > 0$

**then**  $a_{i+1} := a_i, b_{i+1} := t_i$

**else**  $a_{i+1} := t_i, b_{i+1} := b_i$

$$t_{i+1} := \frac{a_{i+1} + b_{i+1}}{2}, i := i + 1$$

**end**

### Theorem

The sequence  $\{t_i\}$  converges to  $t^*$  and  $|t_i - t^*| < (b_0 - a_0)/2^{i+1}$ .

## Gradient method - step size

**Example.**  $\varphi(t) = t^4 + 2t^2 - 3t$ .  $\varphi'(0) = -3$ ,  $\varphi'(1) = 5$ .

$i$	$a_i$	$b_i$	$t_i$	$\varphi'(t_i)$
0	0.000000	1.000000	0.500000	-0.500000
1	0.500000	1.000000	0.750000	1.687500
2	0.500000	0.750000	0.625000	0.476563
3	0.500000	0.625000	0.562500	-0.038086
4	0.562500	0.625000	0.593750	0.212280
5	0.562500	0.593750	0.578125	0.085403
6	0.562500	0.578125	0.570313	0.023241
7	0.562500	0.570313	0.566406	-0.007526
8	0.566406	0.570313	0.568359	0.007831
9	0.566406	0.568359	0.567383	0.000146
10	0.566406	0.567383	0.566895	-0.003692

## Gradient method - step size

Newton method (tangent method): write the first order approximation of  $\varphi'$  at  $t_i$ :

$$\varphi'(t) \simeq \varphi'(t_i) + \varphi''(t_i)(t - t_i)$$

and solve the linear equation  $\varphi'(t_i) + \varphi''(t_i)(t - t_i) = 0$ .

## Newton method

Choose  $t_0 > 0$ , set  $i := 0$

**while**  $\varphi'(t_i) \neq 0$  **do**

$$t_{i+1} := t_i - \frac{\varphi'(t_i)}{\varphi''(t_i)}, i := i + 1$$

**end**

## Theorem

If  $\varphi''(t^*) \neq 0$ , then there exist  $\delta > 0$  and  $C > 0$  such that for any

$t_0 \in (t^* - \delta, t^* + \delta)$  the sequence  $\{t_i\}$  converges to  $t^*$  and

$$|t_{i+1} - t^*| \leq C|t_i - t^*|^2.$$



## Gradient method - step size

**Example.**  $\varphi(t) = t^4 + 2t^2 - 3t$ ,  $t_0 = 1$ .

$i$	$t_i$	$\varphi'(t_i)$
0	1.0000000	5.0000000
1	0.6875000	1.0498047
2	0.5789580	0.0920812
3	0.5674799	0.0009093
4	0.5673642	0.0000001

## Gradient method - zig-zag behaviour

Two subsequent directions are orthogonal:  $(d^k)^T d^{k+1} = 0$

### Theorem

If  $f(x) = \frac{1}{2} x^T Q x + c^T x$ , with  $Q$  positive definite matrix, then

$$\|x^{k+1} - x^*\|_Q \leq \left( \frac{\frac{\lambda_n}{\lambda_1} - 1}{\frac{\lambda_n}{\lambda_1} + 1} \right) \|x^k - x^*\|_Q,$$

where  $\|x\|_Q = \sqrt{x^T Q x}$  and  $0 < \lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_n$  are the eigenvalues of  $Q$ .

## Gradient method - zig-zag behaviour


**Example.**  $f(x) = x_1^2 + 10x_2^2$ , global minimum is  $x^* = (0, 0)$ .

If the starting point is  $x^0 = (10, 1)$ , then

$$x^k = \left( 10 \left( \frac{9}{11} \right)^k, \left( -\frac{9}{11} \right)^k \right), \quad \forall k \geq 0,$$

hence

$$\|x^{k+1} - x^*\| = \frac{9}{11} \|x^k - x^*\| \quad \forall k \geq 0.$$



## Gradient method - Armijo inexact line search

### Gradient method with the Armijo inexact line search

Set  $\alpha, \gamma \in (0, 1)$ ,  $\bar{t} > 0$ . Choose  $x^0 \in \mathbb{R}^n$ , set  $k = 0$ .

```
while  $\nabla f(x^k) \neq 0$  do  
     $d^k = -\nabla f(x^k)$   
     $t_k = \bar{t}$   
    while  $f(x^k + t_k d^k) > f(x^k) + \alpha t_k (d^k)^T \nabla f(x^k)$  do  
         $t_k = \gamma t_k$   
    end  
     $x^{k+1} = x^k + t_k d^k$ ,  $k = k + 1$   
end
```

### Theorem

If  $f$  is coercive, then for any starting point  $x^0$  the generated sequence  $\{x^k\}$  is bounded and any of its cluster points is a stationary point of  $f$ .

## Gradient method - Armijo inexact line search

**Example.** Let  $f(x_1, x_2) = x_1^4 + x_1^2 + x_2^2$ . Set  $\alpha = 10^{-4}$ ,  $\gamma = 0.5$ ,  $\bar{t} = 1$ , choose  $x^0 = (1, 1)$ .

$$d^0 = -\nabla f(x^0) = (-6, -2).$$

Line search. If  $t_0 = 1$  then

$$f(x^0 + t_0 d^0) = 651 > f(x^0) + \alpha t_0 (d^0)^T \nabla f(x^0) = 2.996,$$

if  $t_0 = 0.5$  then

$$f(x^0 + t_0 d^0) = 20 > f(x^0) + \alpha t_0 (d^0)^T \nabla f(x^0) = 2.998,$$

if  $t_0 = 0.25$  then

$$f(x^0 + t_0 d^0) = 0.5625 < f(x^0) + \alpha t_0 (d^0)^T \nabla f(x^0) = 2.999$$

hence the step size is  $t_0 = 0.25$  and the new iterate is

$$x^1 = x^0 + t_0 d^0 = (1, 1) + \frac{1}{4} (-6, -2) = \left(-\frac{1}{2}, \frac{1}{2}\right).$$

## Gradient method - Armijo inexact line search

**Exercise 6.** Solve the problem

$$\begin{cases} \min & 2x_1^4 + 3x_2^4 + 2x_1^2 + 4x_2^2 + x_1x_2 - 3x_1 - 2x_2 \\ & x \in \mathbb{R}^2 \end{cases}$$

by means of the gradient method with the Armijo inexact line search setting  $\alpha = 0.1$ ,  $\gamma = 0.9$ ,  $\bar{\epsilon} = 1$  and starting from the point  $(0, 0)$ .

[Use  $\|\nabla f(x)\| < 10^{-3}$  as stopping criterion.]

## Conjugate gradient method

Search direction involves directions computed at previous iterations.

First, consider the quadratic case:

$$f(x) = \frac{1}{2} x^T Q x + c^T x,$$

where  $Q$  is positive definite. Set  $g = \nabla f(x) = Qx + c$ .

At iteration  $k$ , the search direction is

$$d^k = \begin{cases} -g^0 & \text{if } k = 0, \\ -g^k + \beta_k d^{k-1} & \text{if } k \geq 1, \end{cases}$$

where  $\beta_k$  is such that  $d^k$  and  $d^{k-1}$  are conjugate with respect to  $Q$ , i.e.

$$(d^k)^T Q d^{k-1} = 0.$$

## Conjugate gradient method

- ▶ it easy to compute  $\beta_k$ :

$$\beta_k = \frac{(g^k)^\top Q d^{k-1}}{(d^{k-1})^\top Q d^{k-1}}$$

- ▶ if we perform exact line search, then  $d^k$  is a descent direction
- ▶ the step size given by exact line search is  $t_k = -\frac{(g^k)^\top d^k}{(d^k)^\top Q d^k}$

## Conjugate gradient method (quadratic functions)

Choose  $x^0 \in \mathbb{R}^n$ , set  $g^0 = Q x^0 + c$ ,  $k := 0$

**while**  $g^k \neq 0$  **do**

**if**  $k = 0$  **then**  $d^k = -g^k$

**else**  $\beta_k = \frac{(g^k)^\top Q d^{k-1}}{(d^{k-1})^\top Q d^{k-1}}$ ,  $d^k = -g^k + \beta_k d^{k-1}$

**end**

$t_k = -\frac{(g^k)^\top d^k}{(d^k)^\top Q d^k}$

$x^{k+1} = x^k + t_k d^k$ ,  $g^{k+1} = Q x^{k+1} + c$ ,  $k = k + 1$

**end**



## Conjugate gradient method

**Example.**  $f(x) = x_1^2 + 10x_2^2$ , starting point  $x^0 = (10, 1)$ .

$$k = 0: g^0 = (20, 20), d^0 = -g^0 = (-20, -20),$$

$$t_0 = -((g^0)^T d^0) / ((d^0)^T Q d^0) = 1/11, \text{ hence } x^1 = x^0 + t_0 d^0 = (90/11, -9/11)$$

$$k = 1: g^1 = (180/11, -180/11), \beta_1 = ((g^1)^T Q d^0) / ((d^0)^T Q d^0) = 81/121,$$

$$d^1 = -g^1 + \beta_1 d^0 = (-3600/121, 360/121),$$

$t_1 = -((g^1)^T d^1) / ((d^1)^T Q d^1) = 11/40$ , hence  $x^2 = x^1 + t_1 d^1 = (0, 0)$  which is the global minimum of  $f$ .

## Conjugate gradient method

### Proposition

- ▶ An alternative formula for the step size is  $t_k = \frac{\|g^k\|^2}{(d^k)^T Q d^k}$
- ▶ An alternative formula for  $\beta_k$  is  $\beta_k = \frac{\|g^k\|^2}{\|g^{k-1}\|^2}$
- ▶ If we did not find the global minimum after  $k$  iterations, then the gradients  $\{g^0, g^1, \dots, g^k\}$  are orthogonal
- ▶ If we did not find the global minimum after  $k$  iterations, then the directions  $\{d^0, d^1, \dots, d^k\}$  are conjugate w.r.t.  $Q$  and  $x^k$  is the minimum of  $f$  on  $x^0 + \text{Span}(d^0, d^1, \dots, d^k)$

### Theorem (Convergence)

- ▶ The CG method finds the global minimum in at most  $n$  iterations.
- ▶ If  $Q$  has  $r$  distinct eigenvalues, then CG method finds the global minimum in at most  $r$  iterations.

## Conjugate gradient method

### Theorem (Error bound)

If  $0 < \lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_n$  are the eigenvalues of  $Q$ , then



$$\|x^k - x^*\|_Q \leq 2 \left( \frac{\sqrt{\frac{\lambda_n}{\lambda_1}} - 1}{\sqrt{\frac{\lambda_n}{\lambda_1}} + 1} \right)^k \|x^0 - x^*\|_Q$$



$$\|x^k - x^*\|_Q \leq \left( \frac{\lambda_{n-k+1} - \lambda_1}{\lambda_{n-k+1} + \lambda_1} \right) \|x^0 - x^*\|_Q.$$

## Conjugate gradient method - Exercises

**Exercise 7.** Implement in MATLAB the conjugate gradient method for solving the problem

$$\begin{cases} \min \frac{1}{2}x^T Qx + c^T x \\ x \in \mathbb{R}^n \end{cases}$$

where  $Q$  is a positive definite matrix.

**Exercise 8.** Run the conjugate gradient method for solving the problem

$$\begin{cases} \min 3x_1^2 + 3x_2^2 + 3x_3^2 + 3x_4^2 - 4x_1x_3 - 4x_2x_4 + x_1 - x_2 + 2x_3 - 3x_4 \\ x \in \mathbb{R}^4 \end{cases}$$

starting from the point  $(0, 0, 0, 0)$ . [Use  $\|\nabla f(x)\| < 10^{-6}$  as stopping criterion.]

**Exercise 9.** Consider the following data fitting problem: given the experimental data in fitting.txt, find the best approximating polynomial of degree 3 w.r.t. the Euclidean norm. Solve the problem by means of the conjugate gradient method starting from  $x_0 = 0$ . [Use  $\|\nabla f(x)\| < 10^{-3}$  as stopping criterion.]

## Conjugate gradient method - nonlinear functions

### Conjugate gradient method (nonlinear functions)

Choose  $x^0 \in \mathbb{R}^n$ , set  $k := 0$

**while**  $\nabla f(x^k) \neq 0$  **do**

**if**  $k = 0$  **then**  $d^k = -\nabla f(x^k)$

**else**  $\beta_k = \frac{\|\nabla f(x^k)\|^2}{\|\nabla f(x^{k-1})\|^2}$ ,  $d^k = -\nabla f(x^k) + \beta_k d^{k-1}$

**end**

    Compute the step size  $t_k$

$x^{k+1} = x^k + t_k d^k$ ,  $k = k + 1$

**end**

## Conjugate gradient method - nonlinear functions

### Proposition

- ▶ If  $t_k$  is computed by exact line search, then  $d^k$  is a descent direction.
- ▶ If  $t_k$  satisfies the following conditions:

$$\begin{aligned} f(x^k + t_k d^k) &\leq f(x^k) + \alpha t_k \nabla f(x^k)^\top d^k, \\ |\nabla f(x^k + t_k d^k)^\top d^k| &\leq -\beta \nabla f(x^k)^\top d^k, \end{aligned} \tag{1}$$

with  $0 < \alpha < \beta < 1/2$ , then  $d^k$  is a descent direction.

### Theorem

If  $f$  is coercive, then the conjugate gradient method, where (1) holds, generates a sequence  $\{x^k\}$  such that

$$\liminf_{k \rightarrow \infty} \|\nabla f(x^k)\| = 0.$$

## Newton method

We want to find a stationary point  $\nabla f(x) = 0$ .

At iteration  $k$ , make a linear approximation of  $\nabla f(x)$  at  $x^k$ , i.e.

$$\nabla f(x) \simeq \nabla f(x^k) + \nabla^2 f(x^k)(x - x^k),$$

the new iterate  $x^{k+1}$  is the solution of the linear system

$$\nabla f(x^k) + \nabla^2 f(x^k)(x - x^k) = 0.$$

Note that  $x^{k+1}$  is a stationary point of the quadratic approximation of  $f$  at  $x^k$ :

$$f(x) \simeq f(x^k) + (x - x^k)^T \nabla f(x^k) + \frac{1}{2} (x - x^k)^T \nabla^2 f(x^k) (x - x^k).$$

## Newton method

### Newton method (basic version)

Choose  $x^0 \in \mathbb{R}^n$ , set  $k = 0$

**while**  $\nabla f(x^k) \neq 0$  **do**

Solve the linear system  $\nabla^2 f(x^k)d^k = -\nabla f(x^k)$

$x^{k+1} = x^k + d^k$ ,  $k = k + 1$

**end**

### Theorem (Convergence)

If  $x^*$  is a local minimum of  $f$  and  $\nabla^2 f(x^*)$  is positive definite, then there exists  $\delta > 0$  such that for any  $x^0 \in B(x^*, \delta)$  the generated sequence  $\{x^k\}$  converges to  $x^*$  and

$$\|x^{k+1} - x^*\| \leq C \|x^k - x^*\|^2 \quad \forall k > \bar{k}, \quad (\text{quadratic convergence})$$

for some  $C > 0$  and  $\bar{k} \in \mathbb{N}$ .



## Newton method

**Example.**  $f(x) = 2x_1^4 + 3x_2^4 + 2x_1^2 + 4x_2^2 + x_1x_2 - 3x_1 - 2x_2$  is strongly convex because

$$\nabla^2 f(x) = \begin{pmatrix} 24x_1^2 + 4 & 1 \\ 1 & 36x_2^2 + 8 \end{pmatrix}$$

$k$	$x^k$		$\ \nabla f(x^k)\ $
0	10.000000	5.000000	8189.6317378
1	6.655450	3.298838	2429.6437291
2	4.421132	2.149158	721.6330686
3	2.925965	1.361690	214.6381594
4	1.923841	0.811659	63.7752575
5	1.255001	0.428109	18.6170045
6	0.823359	0.209601	5.0058040
7	0.580141	0.171251	1.0538969
8	0.492175	0.179815	0.1022945
9	0.481639	0.180914	0.0013018
10	0.481502	0.180928	0.0000002

## Newton method

Drawbacks of Newton method:

- ▶ at each iteration we need to compute both the gradient  $\nabla f(x^k)$  and the hessian matrix  $\nabla^2 f(x^k)$
- ▶ local convergence: if  $x^0$  is too far from the optimum  $x^*$ , then the generated sequence can be not convergent to  $x^*$

**Example.** Let  $f(x) = -\frac{1}{16}x^4 + \frac{5}{8}x^2$ .

Then  $f'(x) = -\frac{1}{4}x^3 + \frac{5}{4}x$  and  $f''(x) = -\frac{3}{4}x^2 + \frac{5}{4}$ .

$x^* = 0$  is a local minimum of  $f$  with  $f''(x^*) = 5/4 > 0$ .

The sequence does not converge to  $x^*$  if the method starts from  $x^0 = 1$ :

$x^1 = -1, x^2 = 1, x^3 = -1, \dots$

## Newton method with line search

If  $f$  is strongly convex, we get **global convergence** because  $d^k$  is a descent direction:  $\nabla f(x^k)^\top d^k = -\nabla f(x^k)^\top [\nabla^2 f(x^k)]^{-1} \nabla f(x^k) < 0$ .

## Newton method with line search

Set  $\alpha, \gamma \in (0, 1)$ ,  $\bar{t} > 0$ . Choose  $x^0 \in \mathbb{R}^n$ , set  $k = 0$

**while**  $\nabla f(x^k) \neq 0$  **do**

Solve the linear system  $\nabla^2 f(x^k) d^k = -\nabla f(x^k)$

$t_k = \bar{t}$

**while**  $f(x^k + t_k d^k) > f(x^k) + \alpha t_k (d^k)^\top \nabla f(x^k)$  **do**

$t_k = \gamma t_k$

**end**

$x^{k+1} = x^k + t_k d^k$ ,  $k = k + 1$

**end**

## Theorem (Convergence)

If  $f$  is strongly convex, then for any starting point  $x^0 \in \mathbb{R}^n$  the sequence  $\{x^k\}$  converges to the global minimum of  $f$ . Moreover, if  $\alpha \in (0, 1/2)$  and  $\bar{t} = 1$  then the convergence is quadratic.

## Newton method with line search

**Exercise 10.** Solve the problem

$$\begin{cases} \min & 2x_1^4 + 3x_2^4 + 2x_1^2 + 4x_2^2 + x_1x_2 - 3x_1 - 2x_2 \\ & x \in \mathbb{R}^2 \end{cases}$$

by means of the Newton method with line search setting  $\alpha = 0.1$ ,  $\gamma = 0.9$ ,  $\bar{t} = 1$  and starting from the point  $(0, 0)$ .

[Use  $\|\nabla f(x)\| < 10^{-3}$  as stopping criterion.]

## Quasi-Newton methods

At each iteration  $[\nabla^2 f(x^k)]^{-1}$  is approximated by a positive definite matrix  $H_k$

### Quasi-Newton method

Choose  $x^0 \in \mathbb{R}^n$ , a positive definite matrix  $H_0$ ,  $k = 0$

```
while  $\nabla f(x^k) \neq 0$  do  
     $d^k = -H_k \nabla f(x^k)$   
    Compute step size  $t_k$   
     $x^{k+1} = x^k + t_k d^k$ , update  $H_{k+1}$ ,  $k = k + 1$   
end
```

How to update matrix  $H_k$ ?

## Quasi-Newton methods

$$\nabla f(x^k) \simeq \nabla f(x^{k+1}) + \nabla^2 f(x^{k+1})(x^k - x^{k+1}).$$

Set  $p^k = x^{k+1} - x^k$  and  $g^k = \nabla f(x^{k+1}) - \nabla f(x^k)$ , then

$$\nabla^2 f(x^{k+1}) p^k \simeq g^k, \text{ i.e. } [\nabla^2 f(x^{k+1})]^{-1} g^k \simeq p^k.$$

We choose  $H_{k+1}$  such that

$$H_{k+1} g^k = p^k.$$

Davidon-Fletcher-Powell (DFP) method:

$$H_{k+1} = H_k + \frac{p^k (p^k)^\top}{(p^k)^\top g^k} - \frac{H_k g^k (g^k)^\top H_k}{(g^k)^\top H_k g^k},$$

## Quasi-Newton methods

Another approach: find a matrix  $B_k = (H_k)^{-1}$  approximating  $\nabla^2 f(x^k)$ .

Since  $\nabla^2 f(x^{k+1}) p^k \simeq g^k$ , we impose that  $B_{k+1} p^k = g^k$

Update  $B_k$  as

$$B_{k+1} = B_k + \frac{g^k (g^k)^\top}{(p^k)^\top g^k} - \frac{B_k p^k (p^k)^\top B_k}{(p^k)^\top B_k p^k},$$

hence

$$H_{k+1} = H_k + \left( 1 + \frac{(g^k)^\top H_k g^k}{(p^k)^\top g^k} \right) \frac{p^k (p^k)^\top}{(p^k)^\top g^k} - \frac{p^k (g^k)^\top H_k + H_k g^k (p^k)^\top}{(p^k)^\top g^k}.$$

(Broyden–Fletcher–Goldfarb–Shanno (BFGS) method).

## Derivative-free methods

There are situations where derivatives of the objective function do not exist or are computationally expensive.

**Derivative-free methods** sample the objective function at a finite number of points at each iteration, without any explicit or implicit derivative approximation.

### Definition

A **positive basis** is a set of vectors  $\{v^1, \dots, v^p\} \subset \mathbb{R}^n$  such that:

- ▶ any  $x \in \mathbb{R}^n$  is a **conic** combination of  $v^1, \dots, v^p$ , i.e., there exist

$$\alpha_1, \dots, \alpha_p \geq 0 \text{ such that } x = \sum_{i=1}^p \alpha_i v^i$$

- ▶ for any  $i = 1, \dots, p$ ,  $v^i$  is not a **conic** combination of others  $v^1, \dots, v^p$ .

**Examples:**  $\{e_1, \dots, e_n, -e_1, \dots, -e_n\}$  is a positive basis of  $\mathbb{R}^n$ ;  
 $\{(1, 0), (0, 1), (-1, -1)\}$  is a positive basis of  $\mathbb{R}^2$ .

**Proposition.** If  $\{v^1, \dots, v^p\}$  is a positive basis, then for any  $w \in \mathbb{R}^n \setminus \{0\}$  there is  $i \in \{1, \dots, p\}$  such that  $w^T v^i < 0$ .



## Directional direct-search method

### Directional direct-search method

Choose starting point  $x^0 \in \mathbb{R}^n$ , step size  $t_0 > 0$ ,  $\beta \in (0, 1)$ , tolerance  $\varepsilon > 0$  and a positive basis  $D$ . Set  $k = 0$ .

**while**  $t_k > \varepsilon$  **do**

Order the poll set  $\{x^k + t_k d, \quad d \in D\}$

Evaluate  $f$  at the poll points following the chosen order

**If** there is a poll point s.t.  $f(x^k + t_k d) < f(x^k)$

**then**  $x^{k+1} = x^k + t_k d$ ,  $t_{k+1} = t_k$  (successful iteration)

**else**  $x^{k+1} = x^k$ ,  $t_{k+1} = \beta t_k$  (step size reduction)

**end**

$k = k + 1$

**end**

The method is called coordinate-search method if  $D = \{e_1, \dots, e_n, -e_1, \dots, -e_n\}$ .

## Directional direct-search method

### Theorem

Assume that all the vectors of the positive basis  $D$  are in  $\mathbb{Z}^n$ . If  $f$  is coercive and continuously differentiable, then the generated sequence  $\{x^k\}$  has a cluster point  $x^*$  such that  $\nabla f(x^*) = 0$ .

**Remark 1.** The assumption that vectors of  $D$  are in  $\mathbb{Z}^n$  can be deleted if we accept new iterates which satisfy a “sufficient” decrease condition:

$$f(x^{k+1}) \leq f(x^k) - t_k^2.$$

**Remark 2.** If a **complete** poll step is performed, i.e.,

$$f(x^{k+1}) \leq f(x^k + t_k d) \quad \forall d \in D,$$

then **any** cluster point of  $\{x^k\}$  is a stationary point of  $f$  and  $\lim_{k \rightarrow \infty} \|\nabla f(x^k)\| = 0$ .

## Variants of the Directional direct-search method

### Directional direct-search method

Choose starting point  $x^0 \in \mathbb{R}^n$ , step size  $t_0 > 0$ ,  $\beta \in (0, 1)$ , tolerance  $\varepsilon > 0$ ,  $\gamma \geq 1$  and a set of positive bases  $\mathcal{D}$ . Set  $k = 0$ .

**while**  $t_k > \varepsilon$  **do**

Choose a positive basis  $D \in \mathcal{D}$

Order the poll set  $\{x^k + t_k d, \quad d \in D\}$

Evaluate  $f$  at the poll points following the chosen order

**If** there is a poll point s.t.  $f(x^k + t_k d) < f(x^k)$

**then**  $x^{k+1} = x^k + t_k d$ ,  $t_{k+1} = \gamma t_k$  (successful iteration)

**else**  $x^{k+1} = x^k$ ,  $t_{k+1} = \beta t_k$  (step size reduction)

**end**

$k = k + 1$

**end**

## Directional direct-search method - Exercises

**Exercise 11.** Solve the problem

$$\begin{cases} \min & 2x_1^4 + 3x_2^4 + 2x_1^2 + 4x_2^2 + x_1x_2 - 3x_1 - 2x_2 \\ & x \in \mathbb{R}^2 \end{cases}$$

by means of the directional direct-search method setting  $x^0 = (0, 0)$ ,  $t_0 = 5$ ,  $\beta = 0.5$ ,  $\varepsilon = 10^{-5}$  and the positive basis  $D = \{(1, 0), (0, 1), (-1, 0), (0, -1)\}$ .

**Exercise 12.** Solve the previous problem by means of the directional direct-search method setting  $x^0 = (0, 0)$ ,  $t_0 = 5$ ,  $\beta = 0.5$ ,  $\varepsilon = 10^{-5}$  and the positive basis  $D = \{(1, 0), (0, 1), (-1, -1)\}$ .